

Dissipative Dynamics of Matter Wave Soliton in Nonlinear Optical Lattice

F.Kh. Abdullaev¹, A. Gammal², H.L.F. da Luz² and Lauro Tomio^{1*}

¹ *Instituto de Física Teórica, Universidade Estadual Paulista, 01405-900, São Paulo, Brazil*

² *Instituto de Física, Universidade de São Paulo, 05315-970, São Paulo, Brazil*

(Dated: 01/July/07)

Dynamics and stability of solitons in two-dimensional (2D) Bose-Einstein condensates (BEC), with low-dimensional (1D) conservative plus dissipative nonlinear optical lattices are investigated. In the case of focusing media (with attractive atomic systems) the collapse of the wave packet is arrested by the dissipative periodic nonlinearity. The adiabatic variation of the background scattering length leads to metastable matter-wave solitons. When the atom feeding mechanism is used, a dissipative soliton can exist in focusing 2D media with 1D periodic nonlinearity. In the defocusing media (repulsive BEC case) with harmonic trap in one dimension and one dimensional nonlinear optical lattice in other direction, the stable soliton can exist. This prediction of variational approach is confirmed by the full numerical simulation of 2D Gross-Pitaevskii equation.

PACS numbers: 03.75.Lm;03.75.-b;05.30.Jp

I. INTRODUCTION

The dynamics of optical and matter wave solitons with different type of management of system parameters has been under intensive investigations in the last years [1, 2]. Two types of modulations have been considered: dispersion and nonlinearity management, which can both occur in time and space. Temporal strong and rapid modulations of the dispersion are more interesting in optical fibers due to many advantages of dispersion managed solitons for optical communications[3, 4, 5] and storage of information. Temporal modulations of nonlinearity are promising in fiber ring lasers and Bose-Einstein condensates (BEC) [6, 7, 8]. In the latter case, the suppression of collapse, implying in the existence of stable multidimensional solitons in attractive BEC, and the generation of periodic patterns of matter waves, have been predicted [9, 10, 11, 12, 13, 14, 15, 16, 17]. In optics, nonlinearity managed solitons have also been observed, as described in Refs. [18, 19, 20, 21].

Recently, the attention has been devoted also to the periodic spatial management in nonlinear optics and Bose-Einstein condensates. In optical media, the nonlinear Kerr coefficient can be periodically modulated in space, leading to the problem of optical beam in a 2D medium with nonlinearity management. In BEC, the spatial variation of scattering length is possible [22, 23, 24, 25, 26, 27, 28], for example, by using optically induced Feshbach resonance [29, 30]. In elongated condensates new types of localized matter waves packets can exist. In 2D case, the situation is less clear. The study of one-dimensional (1D) nonlinear periodic potential in two-dimensional (2D) non-linear Schrödinger equation (NLSE) shows that broad solitons are unstable. As verified in Ref. [31], narrow solitons centered on the maximum of the lattice potential can be stable, but the

stability region is so narrow that they are physically unstable. Stable gap solitons can exist in the BEC under combination of a linear and nonlinear periodic potentials [32, 33, 34].

However, models considered till now are strongly idealized. In particular, using the optically induced Feshbach resonances we can generate mixture of conservative and dissipative nonlinear optical lattice. In view of that, around the Feshbach resonance, one can observed non-vanishing contributions to the imaginary part of the scattering length.

In this work, after an analysis of a conservative system with nonlinear optical lattice, we will consider the influence of nonlinear dissipation on the dynamics and the stability of solitons. In particular, we note that the role of such kind of dissipation can be crucial for the existence of solitons in multi dimensional nonlinear optical lattices. Such hope is supported by the well known fact that the homogeneous nonlinear dissipation can arrest collapse in the cubic focusing multi-dimensional NLSE [35]. The possibility of existence of dissipative solitons is investigated, considering compression effects and atom feeding.

The organization of the paper is as follows. The model is described in the next section. In Sec. 3, it is investigated the properties of localized states in case of attractive and repulsive 2D condensates in 1D nonlinear optical lattice, with or without harmonic trap in one of the dimensions. In Sec. 4, it is performed an analysis of the evolution of 2D soliton under 1D periodic nonlinearity and dissipation, using the variational approach and by direct numerical simulation of the GP equation.

II. THE MODEL

Recently, it has been suggested to generate nonlinear optical lattices in BEC by two counter propagating laser beams near the optical induced Feshbach resonance [25, 26]. The spatial variation of the optical intensity leads to a spatial periodic variation of the atomic scattering

*Corresponding author (tomio@ift.unesp.br)

length. Such structure can support new types of localized nonlinear states. The GP equation for the wave function $\psi \equiv \psi(x_1, x_2, t)$ has the form

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \left(\frac{\partial^2 \psi}{\partial x_1^2} + \beta \frac{\partial^2 \psi}{\partial x_2^2} \right) - g(x_1, x_2) |\psi|^2 \psi, \quad (1)$$

where

$$g(x_1, x_2) \equiv \frac{g_0}{2} + (g_1 + i g_2) (\cos^2(kx_1) + \delta_0 \cos^2(kx_2)). \quad (2)$$

g_0 is related to the s -wave two-body scattering length a_s , $g_0 \equiv (4\pi\hbar^2/m)a_s$, with $g_0 > 0$ ($g_0 < 0$) for attractive (repulsive) condensates; g_1 (> 0) is related to the optical intensity; and g_2 parametrize dissipative effects.

The optically induced scattering length and the inelastic collision rate coefficient K_{inel} (imaginary part of a_s) are described by [29]

$$\Re(a_s) = a_{s0} + \frac{1}{2k_i} \left[\frac{\Gamma_{stim}(x)\Delta}{\Delta^2 + (\Gamma_{spon}/2)^2} \right], \quad (3)$$

$$K_{inel} \equiv \Im(a_s) = \frac{2\pi\hbar}{m} \frac{1}{k_i} \left[\frac{\Gamma_{stim}(x)\Gamma_{spon}}{\Delta^2 + (\Gamma_{spon}/2)^2} \right], \quad (4)$$

where a_{s0} is the scattering length without light, Δ is the detuning from the photo-associated resonance, and $\hbar k_i$ is the relative momentum of the collision. Γ_{stim} is the resonant transition rate between the continuum state and the molecular state, proportional to the laser intensity $I(x)$. Far from the resonance, the imaginary part of the scattering length is small, such that $\Im(a) \ll \Re(a)$. It was shown in Ref. [30] that, in the experiment with ^{87}Rb one can obtain optically induced large variations of the scattering length. The laser intensity was 460 W/cm^2 and the variations a_s occurred from $10a_0$ to $190a_0$ (with $a_{s0} = 100a_0$ and a_0 the Bohr radius).

By considering the following variable changes and definitions in (1),

$$\kappa x = 2kx_1, \quad \kappa y = 2kx_2, \quad \tau = \frac{4w_R t}{\kappa^2}, \quad (5)$$

$$w_R = \frac{E_R}{\hbar}, \quad E_R = \frac{\hbar^2 k^2}{2m}, \quad \gamma_{i=0,1,2} = \frac{g_i}{2|g_0|}, \quad (6)$$

we obtain the dimensionless equation

$$i \frac{\partial u}{\partial \tau} + \frac{\partial^2 u}{\partial x^2} + \beta \frac{\partial^2 u}{\partial y^2} + \gamma(x, y) |u|^2 u = 0, \quad (7)$$

where

$$\gamma(x, y) \equiv \gamma_0 + (\gamma_1 + i\gamma_2)(1 + \delta_0 + \cos(\kappa x) + \delta_0 \cos(\kappa y)), \quad (8)$$

and the wave-function was redefined such that

$$u \equiv u(x, y, \tau) = \sqrt{\frac{\kappa^2 |g_0|}{4E_R}} \psi. \quad (9)$$

From (6) to (9), we should note that γ_0 is fixed to $-1/2$ for attractive condensates; and $+1/2$ for repulsive condensates. Different cases can be considered: 1D geometry is realized when $\beta = \delta_0 = 0$. Anisotropic 2D case

is realized for $\beta = 1$, $\delta_0 = 0$. And the 2D isotropic case can be achieved with $\beta = \delta_0 = 1$. Next, we consider more explicitly in our study the anisotropic 2D case, with $\beta = 1$ and $\delta_0 = 0$. As the soliton is completely free in the y -direction, we also examine the possibility to have a harmonic trap $m\omega_y^2 x_2^2/2$. Following the transformations (5), a dimensionless frequency ω is also defined, such that

$$\omega \equiv \kappa^2 \frac{\omega_2}{8w_R}, \quad \frac{m}{2} \omega_2^2 x_2^2 = \left(\frac{4E_R}{\kappa^2} \right) \omega^2 y^2. \quad (10)$$

In order to extend our study of the stability conditions in a few realistic cases, it is also verified the effect of a compression, which can be achieved by an adiabatic time variation of the background value of a scattering length [22], by modifying γ_0 as

$$\gamma_0 \rightarrow \gamma_0(\tau) = \gamma_0 \exp[2\alpha(\tau - \tau_c)] \theta(\tau - \tau_c). \quad (11)$$

Compression effect can also be achieved by a feeding process, which can be described by an additional term $i\alpha_f u$ in the GP equation [36]. Note: If the modulation of nonlinearity in time is induced by increasing the transverse frequency of the trap, then we should multiply the full nonlinear term by $\exp(2\alpha\tau)$. With these considerations, the Eq. (7) can be written as

$$i \frac{\partial u}{\partial \tau} = -\frac{\partial^2 u}{\partial x^2} - \frac{\partial^2 u}{\partial y^2} - \gamma(x, y, \tau) |u|^2 u + \omega^2 y^2 u + i\alpha_f u, \quad (12)$$

where

$$\gamma(x, y, \tau) \equiv \gamma_0(\tau) + (\gamma_1 + i\gamma_2) [1 + \cos(\kappa x)]. \quad (13)$$

In the above, one should take $\alpha_f = 0$ when $\alpha \neq 0$ in (11), as such parameters have similar role in the formalism.

III. CONSERVATIVE SYSTEM

It is useful to describe shortly the solitons and their stability in the conservative case ($\gamma_2 = 0$). One dimensional conservative case has been considered by using a variational approach (VA) in [26]. Using an exact approach, the 2D case with 1D nonlinear optical lattice was studied in [31], where the authors have considered the case with attractive background nonlinearity ($\gamma_1 > 0$). Looking for perspective applications to BEC, we will consider here the 2D problem with 1D nonlinear optical lattice. Following Ref. [26], we start our analysis using the VA formalism.

With $u(x, y, \tau) \equiv v(x, y) \exp(-i\mu\tau)$ in (12), and taking α_f , α and γ_2 equal zero, we obtain

$$\mu v = -\frac{\partial^2 v}{\partial x^2} - \frac{\partial^2 v}{\partial y^2} - (\tilde{\gamma}_0 + \gamma_1 \cos(\kappa x)) v^3 + \omega^2 y^2 v, \quad (14)$$

where we are redefining γ_0 to $\tilde{\gamma}_0 \equiv \gamma_0 + \gamma_1$. In view of our definitions in (5), this implies that for attractive condensed systems we have $\tilde{\gamma}_0 = \gamma_1 + 1/2$; and, for

repulsive ones, $\tilde{\gamma}_0 = \gamma_1 - 1/2$. The sign of $\tilde{\gamma}_0$ gives the sign of the background field. But, we should note that we can have situations where the same $\tilde{\gamma}_0 \geq 0$ can refer to attractive or repulsive condensates. Example:

$\tilde{\gamma}_0 = 1$ with $\gamma_1 = 1/2$ and $\gamma_0 = 1/2$ (attractive);
 $\tilde{\gamma}_0 = 1$ with $\gamma_1 = 3/2$ and $\gamma_0 = -1/2$ (repulsive).

These two situations differ in (14), because the strength of the oscillatory term is different. However, as the results are similar, we prefer to analyze separately the cases of repulsive condensates with negative background field, which occur when $0 < \gamma_1 < 1/2$ ($0 > \tilde{\gamma}_0 \geq -1/2$).

The corresponding averaged Lagrangian L is obtained from the density \mathcal{L} , as

$$L = \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy \mathcal{L}, \quad (15)$$

$$2\mathcal{L} = \mu v^2 - \left| \frac{\partial v}{\partial x} \right|^2 - \left| \frac{\partial v}{\partial y} \right|^2 + \frac{\tilde{\gamma}_0 + \gamma_1 \cos(\kappa x)}{2} v^4 - \omega^2 y^2 v^2. \quad (16)$$

Here, it is interesting to observe that a scaling given by κ is applied to the observables obtained from the above equations, as the root-mean-square radius in x and y directions, chemical potentials, frequencies and energies. In order to see that, we can redefine all the observables using the variable transformation, $\bar{x} \equiv \kappa x$ and $\bar{y} \equiv \kappa y$, such that we have no κ dependence in a new set of observables (represented with a “bar”) that are being calculated. This scaling essentially implies to consider $\kappa \equiv 1$ in all the equations. At the end, the physical observables will be given by the relations (5) (with $\kappa = 1$). For example, in the case of mean-square radius we will have

$$\langle x_1^2 \rangle = \frac{\langle x^2 \rangle}{4k^2} \quad \text{and} \quad \langle x_2^2 \rangle = \frac{\langle y^2 \rangle}{4k^2}. \quad (17)$$

Next, in our VA we consider the Gaussian ansatz

$$v(x, y) = A \exp \left(-\frac{x^2}{2a_1^2} - \frac{y^2}{2a_2^2} \right), \quad (18)$$

which has the normalization given by $N = \pi a_1 a_2 A^2$. The corresponding averaged Lagrangian is given by

$$L = \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy \mathcal{L} = \frac{N}{2} \left[\mu - \left(\frac{1}{2a_1^2} + \frac{1}{2a_2^2} \right) - \frac{\omega^2 a_2^2}{2} + \frac{N}{4\pi a_1 a_2} \left(\tilde{\gamma}_0 + \gamma_1 e^{-\kappa^2 a_1^2/8} \right) \right]. \quad (19)$$

From the Euler-Lagrange equations for the parameters, $\partial L / \partial N = 0$ and $\partial L / \partial a_{i=1,2} = 0$, we obtain:

$$2\mu = \frac{1}{a_1^2} + \frac{1}{a_2^2} - \frac{N}{\pi a_1 a_2} \left(\tilde{\gamma}_0 + \gamma_1 e^{-\frac{\kappa^2 a_1^2}{8}} \right) + \omega^2 a_2^2, \quad (20)$$

$$N = \frac{4\pi a_2}{a_1 [\tilde{\gamma}_0 + \gamma_1 e^{-\kappa^2 a_1^2/8} (1 + \frac{\kappa^2 a_1^2}{4})]}, \quad (21)$$

$$\omega^2 a_2^4 + \frac{a_2^2}{a_1^2} \frac{\left(\tilde{\gamma}_0 + \gamma_1 e^{-\kappa^2 a_1^2/8} \right)}{[\tilde{\gamma}_0 + \gamma_1 e^{-\kappa^2 a_1^2/8} (1 + \frac{\kappa^2 a_1^2}{4})]} - 1 = 0. \quad (22)$$

In the case that $\omega = 0$, this set of equations, for μ , N , and a_2 , can be expressed in terms of a_1 , as

$$\mu_0 = -\frac{1}{a_1^2} \left(\frac{\tilde{\gamma}_0 + \gamma_1 e^{-\kappa^2 a_1^2/8} \left[1 - \frac{\kappa^2 a_1^2}{8} \right]}{\tilde{\gamma}_0 + \gamma_1 e^{-\kappa^2 a_1^2/8} \left[1 + \frac{\kappa^2 a_1^2}{4} \right]} \right) \quad (23)$$

$$a_{2,0} \equiv a_1 \sqrt{\frac{\tilde{\gamma}_0 + \gamma_1 e^{-\kappa^2 a_1^2/8} (1 + \frac{\kappa^2 a_1^2}{4})}{\tilde{\gamma}_0 + \gamma_1 e^{-\kappa^2 a_1^2/8}}} \quad (24)$$

$$N_0 = \frac{4\pi}{\sqrt{\left[\tilde{\gamma}_0 + \gamma_1 e^{-\kappa^2 a_1^2/8} (1 + \frac{\kappa^2 a_1^2}{4}) \right] \left[\tilde{\gamma}_0 + \gamma_1 e^{-\kappa^2 a_1^2/8} \right]}} \quad (25)$$

For the case that $\omega \neq 0$, the relation for a_2 in terms of a_1 can be obtained from (22) and (24):

$$a_2 = \frac{1}{\omega a_{2,0}} \sqrt{\left[\sqrt{\frac{1}{4} + \omega^2 a_{2,0}^4} - \frac{1}{2} \right]}. \quad (26)$$

Equations (26), (20) and (21) form the set of equations for $\omega \neq 0$.

Next, we consider separately the cases of attractive systems, with $\tilde{\gamma}_0 = \gamma_1 + \frac{1}{2} > 0$, and repulsive ones with $\tilde{\gamma}_0 = \gamma_1 - \frac{1}{2} < 0$.

1. Attractive condensate ($\tilde{\gamma}_0 = \gamma_1 + 1/2$)

This case, which corresponds to $\gamma_0 = 1/2$ and $\gamma_1 > 0$, has been investigated recently in [31]. With $\omega = 0$, it is applied the set of equations (23), (25) and (24). In the general case with $\omega \neq 0$, we should consider Eqs. (20), (21), and (26).

Let us first verify the analytic limiting cases of the VA expressions, for $\omega = 0$:

$$\begin{aligned} a_{2,0} &\rightarrow a_1, \text{ for } a_1 \ll 1 \text{ and } a_1 \gg 1; \\ \mu_0 &\rightarrow -1/a_1^2, \text{ for } a_1 \ll 1 \text{ and } a_1 \gg 1; \\ N_0 &\rightarrow \frac{4\pi}{\tilde{\gamma}_0 + \gamma_1} = \frac{4\pi}{2\gamma_1 + 1/2}, \text{ for } a_1 = 0; \\ N_0 &\rightarrow \frac{4\pi}{\tilde{\gamma}_0} = \frac{4\pi}{\gamma_1 + 1/2}, \text{ for } a_1 \rightarrow \infty; \end{aligned}$$

Limiting cases of the VA expressions, for $\omega \neq 0$:

$$\begin{aligned} a_2 &\rightarrow \begin{cases} a_1, & \text{for } a_1 \ll 1; \\ 1/\sqrt{\omega}, & \text{for } a_1 \gg 1 \end{cases} \\ \mu &\rightarrow \begin{cases} -1/a_1^2, & \text{for } a_1 \ll 1 \\ -2/a_1^2 + \omega \rightarrow \omega, & \text{for } a_1 \gg 1; \end{cases} \\ N &\rightarrow \frac{4\pi}{\tilde{\gamma}_0 + \gamma_1}, \text{ for } a_1 = 0; \\ N &\rightarrow \frac{4\pi a_2}{\tilde{\gamma}_0 a_1} \rightarrow 0, \text{ for } a_1 \rightarrow \infty. \end{aligned}$$

In Fig. 1, we plot the corresponding results for the chemical potential μ as a function of N (upper frame) and

N as a function of a_1 (lower frame). Numerical solutions to PDE results were done with algorithm presented in Ref. [37]. Considering the Vakhitov-Kolokolov (VK) criterion [38] for the soliton stability, $d\mu/dN < 0$, from the results given in the upper frame of Fig. 1, we note that the solitons are unstable. This result is in agreement with the prediction of Ref. [31]. We note, from the VA results, that in the limit of large a_1 the system has a tendency to stabilize, indicating that with just a small trapping potential we can produce a stable region. This behavior is shown by the VA results given in Fig. 2. The variational approach, besides an expected small quantitative shift, provides a good qualitative picture of the results when compared with full numerical predictions. If one is first concerned with the stability of the system (instead of the quantitative results of the observables), the VA provides a nice and reliable picture.




FIG. 1: Attractive case, with $\tilde{\gamma}_0 = 1$ and $\gamma_1 = 0.5$. Results for the chemical potential μ , as a function of N (upper frame) and N versus a_1 (lower frame), obtained using variational approach (VA) and full numerical calculations. The variational parameter for the width, a_1 , and the root-mean-square radius, $\sqrt{\langle x^2 \rangle}$, are related by $a_1 = \sqrt{2\langle x^2 \rangle}$ [Actually, the physical observables depend on k as given by (5) and (17).

In our VA, when we keep ω fixed (zero or nonzero) and increase the value of γ_1 , we observe that the general picture in respect to stability of the system does not change. This lead us to conclude that we cannot improve the stability of the system by increasing the strength of the lattice periodicity for attractive condensates. In the following, we are going to analyze the cases with $\tilde{\gamma}_0 < 0$.




FIG. 2: Attractive case, with $\tilde{\gamma}_0 = 1$ and $\gamma_1 = 0.5$. VA results for the chemical potential μ , as a function of N (upper frames) and N versus a_1 (lower frame). The results are given for μ near zero, considering three values of the frequency: $\omega = 0$ (solid line), 0.01 (dashed line) and 0.045 (dotted line).

2. Repulsive condensate, with $\tilde{\gamma}_0 < 0$ ($\tilde{\gamma}_0 = \gamma_1 - 1/2$).

We should remind that by repulsive condensate we mean an atomic system where the particles have originally positive two-body scattering length, such that in (2) we have $g_0 < 0$; or $\gamma_0 = -1/2$. So, given γ_1 (parameter of the spatial periodic variation of the atomic scattering length), $\tilde{\gamma}_0 = \gamma_1 - 1/2$. And, if we also consider a negative background, such that $\tilde{\gamma}_0 < 0$, γ_1 will be restricted to the interval $0 < \gamma_1 < 1/2$.

Some other limitations are applied in the parameters, considering that the widths and N must be real positive quantities. The relation between the widths a_2 and a_1 , Eq. (24) for $\omega = 0$, implies in a limitation to the values of a_1 :

$$e^{-\frac{\kappa^2 a_1^2}{8}} \geq \frac{1}{2\gamma_1} - 1, \rightarrow a_{1,max}^2 = \frac{8}{\kappa^2} \ln \left(\frac{\gamma_1}{\frac{1}{2} - \gamma_1} \right). \quad (27)$$

This limit, $a_{1,max}$, is necessary in order to have a_2 and N real and positive quantities for any values of ω . It will also restrict the actual values of the parameter γ_1 to $1/4 < \gamma_1 < 1/2$. The cases with $\gamma_1 > 1/2$ are also allowed, in principle, without upper limit for a_1 . However, such cases will correspond to positive background field, $\tilde{\gamma}_0 > 0$, that have already been considered in the previous subsection.

In view of the above, let us also verify in this case the analytic VA limits. For $\omega = 0$:

$$\begin{aligned} a_{2,0} &\rightarrow \begin{cases} a_1, & \text{for } a_1 < 1; \\ \infty, & \text{for } a_1 = a_{1,max}. \end{cases} \\ \mu_0 &\rightarrow \begin{cases} -1/a_1^2, & \text{for } a_1 < 1; \\ 1/(2a_1^2), & \text{for } a_1 = a_{1,max}. \end{cases} \\ N_0 &\rightarrow \begin{cases} 8\pi/(4\gamma_1 - 1), & \text{for } a_1 = 0; \\ \infty, & \text{for } a_1 = a_{1,max}. \end{cases} \end{aligned}$$

And for $\omega \neq 0$:

$$\begin{aligned} a_2 &\rightarrow \begin{cases} a_1, & \text{for } a_1 < 1; \\ 1/\sqrt{\omega}, & \text{for } a_1 = a_{1,max}. \end{cases} \\ \mu &\rightarrow \begin{cases} -1/a_1^2, & \text{for } a_1 < 1; \\ 1/(2a_1^2) + \omega, & \text{for } a_1 = a_{1,max}. \end{cases} \\ N &\rightarrow \begin{cases} 8\pi/(4\gamma_1 - 1), & \text{for } a_1 = 0; \\ 32\pi/[(1 - 2\gamma_1)\sqrt{\omega}\kappa^2 a_{1,max}^3], & \text{for } a_1 = a_{1,max}. \end{cases} \end{aligned}$$

In Fig. 3, we plot N versus a_1 and the chemical potential μ versus N , for $\tilde{\gamma}_0 = -0.1$ and $\gamma_1 = 0.4$, considering VA and four values of ω (0, 0.07, 0.1, 0.3). In the case of $\omega = 0$, we also include results obtained from exact PDE calculations. Following the VK criterion for stability, $d\mu/dN < 0$, we notice that stable regions start to appear with $\omega \approx 0.1$. With $\omega > 0.3\kappa^2$ the unstable regions almost disappear. However, as one can observe in the lower frame, the width a_1 is quite limited due to the condition (27). The observables μ and a_i depend on the wave parameter k of to the spatial periodic variation of the atomic scattering length through the scaling relations (5) and (6) with $\kappa = 1$. However, contrary to some discussions and conclusions of Ref. [31], specific values of the parameter k cannot affect the conclusions on stability. In such cases of conservative systems, the stability results from combined effects given by the parameters $\tilde{\gamma}_0$, γ_1 and ω . Our main conclusion is that, without the trapping potential (included in the y -direction), taking $\omega = 0$, the optical lattice cannot stabilize the solutions, neither for repulsive nor for attractive condensates.

In order to further check the role of the optical lattice, for the repulsive case we also investigate the case with constant ω and different values of γ_1 . From the results shown in Fig. 3, for $\gamma_1 = 0.4$, we found appropriate to consider $\omega = 0.07$, which has a marginal stability near $\mu \approx 0.05$. The results are shown in Fig. 4, where we first observe that a larger γ_1 can help to allow the width a_1 to increase, within the limiting condition (27). However, the marginal stability remains for corresponding different values of the chemical potential. In order to keep the plots of Fig.4 for different values of γ_1 in the same frames, we have normalize the number N such that it is equal to one when a_1 is zero.

The plots of the evolution of profiles are presented for $\omega = 0$ and 0.3 in Fig. 5, confirming the VK prediction. The results using the VA have good agreement with PDE prediction, as shown for $\omega = 0$ and 0.3. The profile at $t = 50$ is practically undistinguishable from the initial soliton form.




FIG. 3: Repulsive case, with $\tilde{\gamma}_0 = -0.1$ and $\gamma_1 = 0.4$, for μ versus N (upper frame), and N versus a_1 (lower frame). In both frames, we show the results using the variational approach, for $\omega = 0, 0.07, 0.1$ and 0.3 . In the upper frame, the exact PDE results are also shown in two cases: $\omega = 0$ (for which the system is unstable) and $\omega = 0.3$ (for which the system is stable). In this last case, near the region where the VA presents a small unstable branch ($22 < N < 25$), our exact numerical results are shown only for $N > 24$. As observed, the VA is giving a general picture of the exact solutions.

IV. EVOLUTION OF 2D SOLITON UNDER 1D PERIODIC NONLINEARITY AND DISSIPATION

In this section, we will consider the case we have $\gamma_2 \neq 0$ in (12). To study the dynamics of a 2D soliton with 1D periodic nonlinearity and dissipation, we also apply a variational approach and full numerical calculations. In the Gaussian ansatz (18) we should also include parameters related to dissipative effects and initial conditions. For a bright soliton, the ansatz can be taken in the following form:

$$\begin{aligned} u &= A \exp \left[-\frac{(x-x_0)^2}{2a_1^2} - \frac{y^2}{2a_2^2} \right] \times \\ &\times \exp [ib_1(x-x_0)^2 + ib_2y^2 + i\phi(\tau)], \end{aligned} \quad (28)$$

where b_i ($i=1,2$) are related to dissipative effects, with x_0 and the phase ϕ related to initial condition.

The Lagrangian density for Eq.(12) is

$$\mathcal{L}(x, \tau) = \frac{i}{2}(u_\tau u^* - u^*_\tau u) - |u_x|^2 - |u_y|^2 + \frac{\gamma(x)}{2}|u|^4, \quad (29)$$

where $\gamma(x)$ is given by (13). Next, from the ansatz (28),




FIG. 4: VA results for the repulsive case, with $\tilde{\gamma}_0 = -0.1$ and ω fixed to 0.07, considering $\gamma_1 = 0.35, 0.4, 0.45$ and 0.5. In the upper frame we have μ versus $N/N(a_1 = 0)$; and, in the lower frame, $N/N(a_1 = 0)$ versus a_1 .




FIG. 5: The soliton profile in the stable region predicted by the VK criterion for $\omega = 0.3$, $\tilde{\gamma}_0 = -0.1$, $\gamma_1 = 0.4$, $\mu = 0.1$, $N \approx 33$ at the time $t = 50$. Solid line is x-direction, dashed line is y-direction.

we obtain the corresponding averaged Lagrangian:

$$\begin{aligned}
 L &= \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy \mathcal{L} \\
 &= -\frac{\pi}{2} A^2 a_1 a_2 \left[a_1^2 (b_{1\tau} + 4b_1^2) + \frac{1}{a_1^2} + a_2^2 (b_{2\tau} + 4b_2^2) \right. \\
 &\quad \left. + \frac{1}{a_2^2} + 2\phi_\tau - \frac{A^2}{2} (\tilde{\gamma}_0 + \gamma_1 \cos(\kappa x_0) e^{-\kappa^2 a_1^2/8}) \right] \quad (30)
 \end{aligned}$$

The equations for the soliton parameters $\eta_i = [A, a, b, \phi]$ in the VA are derived from (see for example [39])

$$\frac{\partial L}{\partial \eta_i} - \frac{d}{d\tau} \frac{\partial L}{\partial \eta_{i,\tau}} = \int_{-\infty}^{\infty} dx \int_{-\infty}^{\infty} dy \left[R \frac{\partial u^*}{\partial \eta_i} + c.c. \right], \quad (31)$$

where the perturbation term R is

$$R = -i\gamma_2(1 + \cos(\kappa x))|u|^2 u + i\alpha_f u. \quad (32)$$

Here we are taking into account a linear amplification term (α_f) describing the atoms feeding. Finally, from the above, we obtain the following system of five coupled ordinary differential equations (ODE) to be solved for the parameters of our variational approach (VA):

$$(A^2 a_1 a_2)_\tau = -\gamma_2 A^4 a_1 a_2 e^{-\kappa^2 a_1^2/8} \cos(\kappa x_0) - \gamma_2 A^4 a_1 a_2 + 2\alpha_f A^2 a_1 a_2, \quad (33)$$

$$(A^2 a_1^3 a_2)_\tau = 8A^2 a_1^3 a_2 b_1 - \frac{\gamma_2}{2} A^4 a_1^3 a_2 [1 + \frac{1}{4} \cos(\kappa x_0) (4 - \kappa^2 a_1^2) e^{-\kappa^2 a_1^2/8}] + 2\alpha_f A^2 a_1^3 a_2, \quad (34)$$

$$(A^2 a_1 a_2^3)_\tau = 8A^2 a_1 a_2^3 b_2 - \frac{\gamma_2}{2} A^4 a_2^3 a_1 [1 + e^{-\kappa^2 a_1^2/8} \cos(\kappa x_0)] + 2\alpha_f A^2 a_1 a_2^3, \quad (35)$$

$$b_{1\tau} = \frac{1}{a_1^4} - 4b_1^2 - \frac{\tilde{\gamma}_0 A^2}{4a_1^2} - \frac{\gamma_1 A^2}{4a_1^2} \cos(\kappa x_0) e^{-\kappa^2 a_1^2/8} (1 + \frac{\kappa^2 a_1^2}{4}), \quad (36)$$

$$b_{2\tau} = \frac{1}{a_2^4} - 4b_2^2 - \frac{\tilde{\gamma}_0 A^2}{4a_2^2} - \frac{\gamma_1 A^2}{4a_2^2} \cos(\kappa x_0) e^{-\kappa^2 a_2^2/8}. \quad (37)$$

By taking into account that the norm $N = \pi A^2 a_1 a_2$, with $i, j = 1, 2$ ($i \neq j$), we have:

$$N_\tau = -\frac{\gamma_2 N^2}{\pi a_1 a_2} \left[1 + e^{-\kappa^2 a_1^2/8} \cos(\kappa x_0) \right] + 2\alpha_f N, \quad (38)$$

$$(a_i^2)_\tau = 8a_i^2 b_i + \frac{\gamma_2 N a_i}{2\pi a_j} \left[1 + e^{-\kappa^2 a_1^2/8} \cos(\kappa x_0) \left(1 + \delta_{i,1} \frac{\kappa^2 a_1^2}{4} \right) \right] \quad (39)$$

$$b_{i,\tau} = \frac{1}{a_i^4} - 4b_i^2 - \frac{N}{4\pi a_i^3 a_j} \left(\gamma_0 + \gamma_1 \left[1 + e^{-\kappa^2 a_1^2/8} \cos(\kappa x_0) \left(1 + \delta_{i,1} \frac{\kappa^2 a_1^2}{4} \right) \right] \right) \quad (40)$$

In the next, we present some of our results, when considering periodic nonlinearity with dissipative effects. Considering the scaling of observables with κ , discussed for the conservative systems in section III, which can also be verified in the present case, we have the corresponding transformation $b_i \rightarrow b_i/\kappa^2$.

In Fig. 6, we have results for the of full numerical simulations (PDE) for the evolution of the matter wave packet under combination of the conservative and dissipative nonlinear optical lattice in the case of the attractive condensate $\gamma_0 = 1/2$. As we can see the collapse is arrested by the dissipative nonlinear optical lattice. The results are compared with the prediction of the VA approach (ODE). We observe a good agreement of the VA with full-numerical calculations.

We also have investigated the role of a deviation δ of the given norm from the critical norm, in the initial wave packet: $A \rightarrow A(1 + \delta)$. The results of the full numerical simulations are presented in Fig. 7. Increasing the deviation δ from the critical norm, multiple peaks are observed, corresponding to revivals of the wave packet during the collapse. The number of peaks grows as δ varies from 0.02 to 0.5. The focusing-defocusing cycles connect the action of the periodically varying in the space with the inelastic three-body interactions. In linear conservative optical lattice, with inelastic three-body interactions, the focusing-defocusing oscillations have been studied in Ref. [40].

The spreading out of the pulse after the collapse is arrested, observed in Fig. 6, can be compensated, as we note previously, by an adiabatic variation of the background scattering length, described by a time variation of γ_0 . When we consider a time dependent γ_0 , as given in Eq. (11), the feeding term parameter α_f should be




FIG. 6: Results for the amplitude as a function of τ , using variational approach (VA) and full numerical calculations. The VA and the full numerical calculations have the value of μ fixed to the same value for $\tau = 0$, implying in a small shift of $A(0)$, as shown by the results.

zero, because one can show (with a redefinition of the wave-function) that it has a similar effect. In Fig. 8, we show our full numerical results confirming the stabilization of the condensate after the collapse was arrested. The mechanism of this stabilization was given by an appropriate tuning of the parameters α and τ_c of Eq. (11).

V. CONCLUSION

Dynamics and stability of matter-wave solitons in the mixture of conservative and dissipative nonlinear optical lattices are investigated, considering 2D BEC, with 1D conservative plus dissipative nonlinear optical lattices.




FIG. 7: Results for the amplitude as a function of τ , showing the collapsing behavior for $\kappa = 4\pi$. The dissipation is fixed to $\gamma_2 = 0.0025$ and δ is increasing from 0.02 to 0.5.




FIG. 8: Soliton stabilization via compression effect, with $\tau_c = 5$ and $\alpha = 0.018$ in Eq. (11). The other parameter are $\gamma_2 = 0.006$, $\delta = 0.01$, and $\kappa = 4\pi$.

In the first part of this work, it was analyzed conservative systems, with nonlinear optical lattices, for attractive and repulsive condensates. It was clarified the role of the scales when calculating the observables as the chemical potential and the widths. Our conclusion is that, in a 2D system, a nonlinear periodic lattice in one direction by itself cannot give stable solutions, satisfying the VK criterium [38]. Such periodic lattice in the x -direction cannot compensate the collapsing effect which results from the other dimension. We verify that stable solutions can be obtained by controlling the soliton with a harmonic trap in the y -direction. For repulsive condensates, the 2D stable soliton can exist in the geometry with 1D nonlinear optical lattices in one direction and harmonic trap in the other direction.

In the second part of the work we analyze the dynamics of the above 2D system, with periodic nonlinearity in x -direction and without trap in the y -direction, when we add non-conservative nonlinear optical lattice terms. We show that the collapse of the condensate can be arrested by a dissipative periodic nonlinearity. To study the evolution of 2D wavepacket we apply the time-dependent variational approach. To compensate the wavepacket broadening, the adiabatic time variation of scattering length is used. It is shown that the metastable dissipative soliton can exist in 2D condensate with 1D periodic nonlinearity. Analytical predictions are confirmed by the numerical simulations of full 2D GP equation.

Acknowledgments

We thank Fundação de Amparo à Pesquisa do Estado de São Paulo for partial support. LT and AG also thank Conselho Nacional de Desenvolvimento Científico e Tecnológico for partial support.

-
- [1] F.Kh. Abdullaev, A. Gammal, A.M. Kamchatnov and L. Tomio, *Int. J. Mod. Phys. B* **19**, 3415 (2005).
 - [2] B.A. Malomed, *Soliton Management in Periodic Systems* (Springer, New York, 2006).
 - [3] I. Gabitov, and S.K. Turitsyn, *Opt. Lett.* **21**, 327 (1996).
 - [4] N.J. Smith *et al.*, *Electr. Lett.* **32**, 54 (1996).
 - [5] F.Kh. Abdullaev, B.B. Baizakov, and M. Salerno, *Phys. Rev. E* **68**, 066605 (2003).
 - [6] L. Bergé *et al.*, *Opt. Lett.* **25**, 1037 (2000).
 - [7] I. Towers and B.A. Malomed, *J. Opt. Soc. Am. B* **19**, 537 (2002).
 - [8] Z.Y. Xu, Y.V. Kartashov, and L. Torner, *Phys. Rev. Lett.* **95**, 113901 (2005).
 - [9] F.Kh. Abdullaev, A.M. Kamchatnov, V.V. Konotop, and V. Brazhnyi, *Phys. Rev. Lett.* **90**, 230402 (2003).
 - [10] F.Kh. Abdullaev, J.G. Caputo, B.A. Malomed, and R.A. Kraenkel, *Phys. Rev. A* **67**, 013605 (2003).
 - [11] H. Saito and M. Ueda, *Phys. Rev. Lett.* **90**, 040403 (2003).
 - [12] G.D. Montesinos, V.M. Perez-Garcia, and P.J. Torres, *Physica D* **191**, 193 (2004).
 - [13] V. Zharntsky and D. Pelinovsky, *Chaos* **15**, 037105 (2005).
 - [14] F.Kh. Abdullaev, E.N. Tsoy, B.A. Malomed, and R.A. Kraenkel, *Phys. Rev. A* **68**, 053606 (2003).
 - [15] F.Kh. Abdullaev and J. Garnier, *Phys. Rev. E* **72**, 035603(R) (2005).
 - [16] P.G. Kevrekidis, D.E. Pelinovsky, and A. Stefanov, *J. Phys. A* **39**, 479 (2006); M.A. Porter, P.G. Kevrekidis, B.A. Malomed, and D.J. Frantzeskakis, *Physica D* **229**, 104 (2007).
 - [17] S.K. Adhikari, *Phys. Rev. A* **69**, 063613 (2004).
 - [18] Q. Quraishi, S.T. Cundiff, B. Ilan, and M. Ablowitz, *Phys. Rev. Lett.*, **94**, 243904 (2005).
 - [19] M. Centurion, M.A. Porter, P.G. Kevrekidis, and D. Psaltis, *Phys. Rev. Lett.* **97**, 033903 (2006).

- [20] P.J. Torres, Nonlinearity **19**, 2103 (2006).
- [21] A. Ciattoni, C. Rizza, E. DelRe and E. Palange, Phys. Rev. Lett. **98**, 043901 (2007).
- [22] F.Kh. Abdullaev and M. Salerno, J. Phys. B **36**, 2851 (2003).
- [23] F.Kh. Abdullaev, A. Gammal, and L. Tomio, J. Phys. B **37**, 635 (2004).
- [24] G. Theocharis et al. Phys. Rev. A **72**, 033614 (2005).
- [25] F.Kh. Abdullaev and J. Garnier, Phys. Rev. A **72**, 061605(R) (2005).
- [26] H. Sakaguchi and B.A. Malomed, Phys. Rev. E **72**, 046610 (2005); Phys. Rev. E **73**, 026601 (2006).
- [27] J. Garnier and F.Kh. Abdullaev, Phys. Rev. A **74**, 01304 (2006).
- [28] J. Belmonte-Beita, V.M. Perez-Garcia, V. Vekslerchik, and P. Torres, Phys. Rev. Lett. **98**, 064102 (2007).
- [29] P.O. Fedichev, Yu. Kagan, G.V. Schlyapnikov, and J.T.M. Walraven, Phys. Rev. Lett. **77**, 2913 (1996).
- [30] M. Theis, *et al.*, Phys. Rev. Lett. **93**, 123001 (2004).
- [31] G. Fibich, Y. Sivan, and M.I. Weinstein, Physica D **217**, 31 (2006); Phys. Rev. Lett. **97**, 193902 (2006).
- [32] Y. Bludov and V.V. Konotop, Phys. Rev. A **74**, 043616 (2006).
- [33] F.Kh. Abdullaev, A.A. Abdumalikov, and R.M. Galimzyanov, Phys.Lett. A in press (2007).
- [34] G. Dong and B. Hu, Phys. Rev. A **75**, 013625 (2007).
- [35] G. Fibich, SIAM J. Appl. Math. **61**, 1680 (2001).
- [36] P.D. Drummond and K.V. Kheruntsyan, Phys. Rev. A **63**, 013605 (2001).
- [37] M. Brtka, A. Gammal, and L. Tomio, Phys.Lett. A **359**, 339 (2006).
- [38] N. G. Vakhitov and A. A. Kolokolov, Radiophys. Quantum Electron. **16**, 783 (1973).
- [39] V. Filho, F.Kh. Abdullaev, A. Gammal, and L. Tomio, Phys. Rev. A **63**, 053603 (2001).
- [40] F.Kh. Abdullaev and M. Salerno, Phys. Rev. A **72**, 033617 (2005).